

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Test fabric for color fastness. Contains a dilute oxidizing agent. (Always refer to manufacturer's care instructions before using equipment or devices). This product is not for use on medical device surfaces. **FOR USE AS A ONE-STEP HOSPITAL DISINFECTANT:** 1. Pre-clean visibly soiled areas. 2. Apply ready-to-use solution to disinfect hard, non-porous surfaces with mechanical spray device, hand pump or coarse trigger spray device. For spray applications, spray 6 – 8 inches from surface. 3. Ensure treated surface is thoroughly wet. 4. Treated surfaces must remain visibly wet for 60 seconds. For SARS-CoV-2, MRSA, E. coli and Listeria treated surfaces must remain visibly wet for only 15 seconds. 5. Wipe dry with a clean cloth or allow to air dry. 6. Food contact surfaces must be rinsed with a clean potable water. **FOR USE AS SOFT SURFACE SANITIZER:** 1. Pre-clean visibly soiled areas. 2. Apply ready-to-use solution to soft surfaces with a mechanical spray device, hand pump or trigger spray device. For spray applications, spray 6 – 8 inches from surface. 3. Treated surfaces must remain visibly wet for 5 minutes. 4. Food contact surfaces must be rinsed with a clean potable water. Test fabric for color fastness. Contains a dilute oxidizing agent. (Always refer to manufacturer's care instructions before using equipment or devices). This application is for spot treatment only. Fabrics must be immersed for treatment beyond spot treatment. **DEODORIZATION:** This product works by oxidation, not by masking of odors. Eliminates odor causing bacteria caused by pet odors, fecal, urine, vomit, smoke (tobacco, marijuana, fire), musty odors, stale-cooking odors. 2. Simply spray, fog, or wipe at full strength and let air dry. 3. When fogging to deodorize apply 1 gallon per 5000 square feet following fogger manufacturer's directions for use. This product must come into contact with the cause of the odor to be effective. For pet urine stains in carpet, blot urine as dry as possible then saturate stain with this product at full strength to carpet pad. **FOR USE AS CLEANER, DEGREASER, 1:** Prepare a use solution of 1 part product to 20 parts water for cleaning applications. 2. Prepare a use solution of 1 part product to 10 parts water for degreaser applications. 3. Prepare this product at full strength for visibly soiled cleaner and degreaser applications. 4. Prepare a use solution of 1 part product to 10 parts water for cleaning visibly soiled carpets with a carpet extractor. 5. Prepare a use solution of 1 part product to 20 parts water for general carpet cleaning with a carpet extractor. 6. Ensure treated surface is thoroughly wet. 7. Treated surfaces must remain visibly wet for 60 seconds. 8. Wipe dry with a clean cloth or allow to air dry. 9. Food contact surfaces must be rinsed with a clean potable water. 10. To clean nonporous surfaces spray soiled area, then wipe clean. –or– for spot cleaning, spray and wipe clean with damp sponge –or– mop or cloth. 11. To spot clean carpet, spray at full strength on soiled area until wet. Gently blot area with a clean, damp, color safe cloth. Repeat as needed (for stubborn stains). Let air dry. 12. To use as a pre-spray (pre-cleaner) on carpet, spray at full strength evenly coating the soiled area (until wet). Test fabric for color fastness. Contains a dilute oxidizing agent. (Always refer to manufacturer's care instructions before using equipment or devices) **ALLERGEN REMOVAL:** To clean and reduce specified allergens: 1. Set trigger to SPRAY. Spray, wait 1 minute, and wipe excess. 2. Allow to air dry. (If streaking is observed, wipe with a clean, damp cloth or paper towel.) Allergens: Cockroach matter –or– particles, Dust mite matter –or– particles, Pet dander –or– dog and cat dander. **FOR USE AS FUNGICIDE:** Pre-clean visibly soiled areas. 2. Apply ready-to-use solution to hard, non-porous surfaces with a mechanical spray device, hand pump or trigger spray device. For spray applications, spray 6 – 8 inches from surface. 3. Treated surfaces must remain visibly wet for 10 minutes. 4. Food contact surfaces must be rinsed with a clean potable water. **FOR USE TO CONTROL MOLD:** 1. Pre-clean, removing as much surface soil, and mold as possible before use. 2. Apply ready-to-use solution, do not dilute, to hard non-porous surfaces with a mechanical spray device, hand pump or trigger spray device. For spray applications, spray 6 – 8 inches from surface. 3. Treated surfaces must remain visibly wet for 10 minutes. 4. Allow to air dry. 5. Food contact surfaces must be rinsed with a clean potable water. For use on carpet and textiles thoroughly moisten affected areas with ready-to-use solution. After a flood or water damage, vacuum or extract as much water as possible before application. Dry carpet within 24 hours to eliminate conditions favorable to mold growth. To assist the drying process, the use of fans, dehumidifiers, heaters or other methods of ventilation are encouraged. Force dry by use of fan or dehumidifier. After thorough drying, the carpet or textile should be vacuumed thoroughly, and the vacuum cleaner bag discarded in a well-sealed plastic bag or container. If a bagless vacuum is used spray container with undiluted product and let stand for 4 minutes. If carpet or textile is not readily accessible for application of product to obtain thorough wetting, the carpet or textile must be removed and discarded in a manner consistent with application regulations and instructions. Consult with your local county extension office or EPA, (<https://www.epa.gov/mold/moldresources.html>) for more information. If the mold growth extends over an area greater than 10 square feet, consider hiring a professional for remediation. Mold damage caused by clean water may be handled by a non-professional; however, mold damage resulting from contaminated water should be handled by a professional who has experience cleaning buildings damaged by contaminated water. If you have health concerns, consult a health professional prior to mold cleanup and pesticide application. Take precautions to limit your exposure to mold and mold spores. Wear protective equipment, such as a N-95 respirator, to limit your exposure to airborne mold. Gloves and goggles must also be worn to protect your hands and eyes. If you know or suspect mold has contaminated the HVAC&R system, do not operate the system. Operating a HVAC&R system that has been contaminated with mold may cause the mold to spread throughout the building. Prior to the use of a pesticide, inspection and proper cleaning of the components to be treated must be performed. If your state requires licensing or certification of persons who inspect or clean such components, the inspection and cleaning must be done by a licensed or certified person. Removed and discarded in a manner consistent with application regulations and instructions. Please consult with your local county extension office or EPA, (<https://www.epa.gov/mold/moldresources.html>) for more information. **FOR USE AS A FUNGISTAT, MILDEWSTAT OR MILDEWICIDE:** A nonpublic health use for aesthetic purposes to control odor and stain causing bacteria. 1. Pre-clean visibly soiled areas. 2. Apply ready-to-use solution to hard, non-porous surfaces with a mechanical spray device, hand pump or trigger spray device. For spray applications, spray 6 – 8 inches from surface. 3. Treated surfaces must remain visibly wet for 10 minutes. 4. Food contact surfaces must be rinsed with a clean potable water.

SoRite KILLS: Bacteria – Broad Spectrum/Hospital Disinfection (RTU) 60 Sec. Contact Time: *Pseudomonas aeruginosa* (ATCC 15442), *Salmonella enterica* (ATCC 10708), *Staphylococcus aureus* (ATCC 6538) **Bacteria Additional (RTU) 15 Sec. Contact Time:** *Escherichia coli* (ATCC 11229), *Methicillin Resistant Staphylococcus aureus – MRSA* (ATCC 33591), *Listeria monocytogenes* (ATCC 19111), *Streptococcal pneumoniae* (ATCC 51936) **60 Sec. Contact Time:** *Carbapenem-resistant Enterobacteriaceae – CRE* (ATCC BAA-2471), *Klebsiella pneumoniae* (ATCC 4352), *Vancomycin Resistant Enterococcus faecium – VRE* (ATCC 51559) **Bacteria – Virucidal (RTU) 15 Sec. Contact Time:** SARS-Related Coronavirus 2 – SARS-CoV-2 (strain USA-WA1/2020, Source BEI Resources, NR-52281) **60 Sec. Contact Time:** Influenza Swine Flu A Virus (strain A/Swine/1976/31, Source: Charles River Laboratories), Influenza Swine Flu B Virus, Hepatitis B Virus (DHBV as surrogate) (strain Grimaud, HepadnaVirus Testing, Inc.), Hepatitis C Virus (BVDV as surrogate) (strain NADL, Source: American BioResearch Lab), Respiratory syncytial Virus – (RSV) (strain Long, ATCC VR-26), Human immunodeficiency Virus type 1 – (HIV-1) (strain 11IB, Source: ZeptoMetrix) **10 Min. Contact Time:** Murine Norovirus (strain MNV-G, Source: Yale University) **Fungicidal/Mold Remediation-Disinfection (RTU) 10 Min. Contact Time:** *Trichophyton interdigitale* (ATCC 9533) **Bacteria-Hard Non Porous Non-Food Contact Surface Sanitizer 30 Sec. Contact Time:** *Klebsiella aerogenes* (ATCC 6538), *Staphylococcus aureus* (ATCC 6538) **Bacteria-Soft Non-Food Contact Surface Sanitizer (RTU) 5 Min. Contact Time:** *Staphylococcus aureus* (ATCC 6538), *Klebsiella aerogenes* (ATCC 13048) **Bacteria – Broad Spectrum/Hospital Disinfection (Use Dilution Method) 10 Min. Contact Time:** *Pseudomonas aeruginosa* (ATCC 15442), *Salmonella enterica* (ATCC 10708), *Staphylococcus aureus* (ATCC 6538)

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal. **PESTICIDE STORAGE:** Store in original closed container in a cool, dry place away from heat and open flame. Do not allow product to become overheated in storage. Avoid prolonged storage temperature above 53°C, 128°F and below 0°C, 32°F. This may cause increased degradation of product, which will decrease product effectiveness.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Discard excess or used (product) solution in drain with running water.

CONTAINER HANDLING: Non-refillable Container. Do not reuse or refill this container. Triple rinse container promptly after emptying. Triple rinse as follows: Fill the container ¼ full with water and recap. Shake for 10 seconds. Drain for 10 seconds after the flow begins to drip. Follow Pesticide Disposal instructions for rinsate disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

FIRST AID: IF IN EYES: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice. **GENERAL INFORMATION:** Have the product container or label with you when calling a poison control center or doctor or going for treatment. For non-emergency and general information on product use, etc., information pertaining to this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378, Monday – Friday, 8:00 am – 12:00 pm Pacific.

EPA Reg. No. 91603-3
EPA Establishment No.: 10508-OH-001, 10508-TN-001
NOTICE: Seller expressly warrants that PRODUCT conforms to its chemical description. There are no other warranties associated with the sale of PRODUCT.

Manufactured by



1109 Woodland Street #68223,
Nashville, TN 37206
Customer Service:(888) 379-3232
aseptichealth.com
Visit us at aseptichealth.com for SDS



CAGE #8PW71

**Escherichia coli*, MRSA, *Listeria monocytogenes* and *Streptococcal pneumoniae* ^SARS-CoV-2